



PCT/AU03/00388


REC'D 25 APR 2003

WIPO

PCT

Patent Office
Canberra

I, TERESA KOLODZIEJCZYK, TEAM LEADER EXAMINATION SUPPORT
AND SALES hereby certify that annexed is a true copy of the Provisional
specification in connection with Application No. PS 1621 for a patent by
MEDVET SCIENCE PTY. LTD. as filed on 08 April 2002.



WITNESS my hand this
Ninth day of April 2003

TERESA KOLODZIEJCZYK
TEAM LEADER EXAMINATION
SUPPORT AND SALES

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Medvet Science Pty. Ltd.

A U S T R A L I A
Patents Act 1990

PROVISIONAL SPECIFICATION

for the invention entitled:

"A Method of Modulating Cellular Activity - III"

The invention is described in the following statement:

A METHOD OF MODULATING CELLULAR ACTIVITY - III

FIELD OF THE INVENTION

5 The present invention relates generally to a method of modulating cellular activity and to agents for use therein. More particularly, the present invention provides a method of modulating cellular activity by modulating phosphorylation of sphingosine kinase and, thereby, its activation. In a related aspect, the present invention provides a method of modulating sphingosine kinase functional activity via modulation of its phosphorylation
10 and agents for use therein. The present invention still further extends to sphingosine kinase variants and to derivatives, analogues, chemical equivalents and mimetics thereof exhibiting reduced and/or ablated capacity to undergo phosphorylation. The method and molecules of the present invention are useful, *inter alia*, in the treatment and/or prophylaxis of conditions characterised by aberrant, unwanted or otherwise inappropriate
15 cellular and/or sphingosine kinase functional activity. The present invention is further directed to methods for identifying and/or designing agents capable of modulating sphingosine kinase phosphorylation.

BACKGROUND OF THE INVENTION

20

Bibliographic details of the publications referred to by author in this specification are collected at the end of the description.

The reference to any prior art in this specification is not, and should not be taken as, an
25 acknowledgment or any form of suggestion that that prior art forms part of the common general knowledge in Australia.

Sphingosine kinase catalyses the formation of sphingosine 1-phosphate (S1P), a lipid messenger that plays important roles in a wide variety of mammalian cellular processes
30 (Pyne *et al.* 2000; Spiegel 1999). S1P is mitogenic in various cell types and triggers a diverse range of important regulatory pathways including; mobilisation of intracellular

calcium by an inositol triphosphate independent pathway (Mattie *et al.* 1994), activation of phospholipase D (Desai *et al.* 1992), inhibition of c-Jun N-terminal kinase (JNK) (Cuvilliver *et al.* 1998), inhibition of caspases (Cuvilliver *et al.* 1998), adhesion molecule expression (Xia *et al.* 1998), and stimulation of DNA binding activity of NF- κ B (Xia *et al.* 5 2002) and transcription factor activator protein-1 (AP-1) (Su *et al.* 1994).

In addition to its role in cellular proliferation and survival, S1P appears to have other functions in the cell. For example, recent studies have shown that S1P is an obligatory signalling intermediate in adhesion molecule expression of vascular endothelial cells (Xia 10 *et al.* 1998), suggesting a likely role in inflammation and atherosclerosis.

Cellular levels of S1P are largely mediated by the activity of sphingosine kinase, and to a lesser extent by its degradation by S1P lyase (Van Veldhoven *et al.* 2000) and S1P phosphatase (Mandala *et al.* 2000) activities. Basal levels of S1P in the cell are generally 15 low (Spiegel *et al.* 1998), but can increase rapidly and transiently when cells are exposed to various mitogenic agents. This response is a direct consequence of an increase in sphingosine kinase activity in the cytosol and can be prevented by the addition of sphingosine kinase inhibitors. This places sphingosine kinase, and its activation, in a central and obligatory role in mediating the observed effects attributed to S1P in the cell. 20 However, at present almost nothing is known of the mechanism(s) leading to sphingosine kinase activation.

Sphingosine kinase can be very rapidly activated by wide variety of cell agonists. While the response differs between cell types, these stimuli include TNF α (Xia *et al.* 1998); 25 Pitson *et al.* 2000) (Fig 1), platelet-derived growth factor (Olivera *et al.*, 1993), epidermal growth factor (Meyer zu Heringdorf *et al.* 1999), nerve growth factor (Rius *et al.*, 1997), vitamin D3 (Kleuser *et al.* 1998), phorbol esters (Pitson *et al.* 2000; Buehrer *et al.* 1996), acetylcholine (muscarinic agonists) (Meyer zu Heringdorf *et al.* 1998), and crosslinking of the immunoglobulin receptors Fc ϵ R1 (Choi *et al.* 1996) and Fc γ R1 (Melendez *et al.* 1998). 30 In all cases this sphingosine kinase activation increases the V_{\max} of the reaction while leaving the substrate affinities (K_m) unaltered.

- 3 -

The molecular mechanisms that couple these agonists to activation of sphingosine kinase activity remain largely unknown. Accordingly, there is a need to elucidate these mechanisms and to develop methods of regulating cellular activities via regulation of the
5 sphingosine kinase signalling pathways.

In work leading up to the present invention, the inventors have determined that phosphorylation of sphingosine kinase is essential for its activation. Further the inventors have identified the phosphorylation sites on the sphingosine kinase molecule. Further, it
10 has been determined that phosphorylation of sphingosine kinase is performed by a proline-directed protein kinase, more particularly ERK2.

SUMMARY OF THE INVENTION

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will
 5 be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

Specific mutations in amino acid sequence are represented herein as "Xaa₁nXaa₂" where Xaa₁ is the original amino acid residue before mutation, n is the residue number and Xaa₂
 10 is the mutant amino acid. The abbreviation "Xaa" may be the three letter or single letter amino acid code. A mutation in single letter code is represented, for example, by X₁nX₂ where X₁ and X₂ are the same as Xaa₁ and Xaa₂ respectively. In terms of both the mutation and the human sphingosine kinase protein sequence in general, the amino acid residues for human sphingosine kinase are numbered with the residue serine(S) in the
 15 motif KTPASPVVVQ <400>1 being numbered 225.

One aspect of the present invention provides a method of modulating sphingosine kinase functional activity, said method comprising contacting said sphingosine kinase with an effective amount of an agent for a time and under conditions sufficient to modulate
 20 phosphorylation of said sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates sphingosine kinase activity.

Another aspect of the present invention provides a method of modulating human
 25 sphingosine kinase functional activity, said method comprising contacting said human sphingosine kinase with an effective amount of an agent for a time and under conditions sufficient to modulate phosphorylation of said human sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said human sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said
 30 human sphingosine kinase activity.

Yet another aspect of the present invention provides a method of modulating sphingosine kinase functional activity, said method comprising contacting said sphingosine kinase with an effective amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of said sphingosine kinase at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰,
 5 wherein inducing or otherwise agonising said phosphorylation up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said sphingosine kinase activity.

Still another aspect of the present invention is directed to a method of modulating
 10 sphingosine kinase functional activity, said method comprising contacting said sphingosine kinase with an effective amount of an agent for a time and under conditions sufficient to modulate proline-directed protein kinase catalysed sphingosine kinase phosphorylation wherein inducing or otherwise agonising said phosphorylation up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation
 15 down-regulates sphingosine kinase activity.

In a related aspect the present invention provides a method of modulating sphingosine kinase functional activity, said method comprising contacting said sphingosine kinase with an effective amount of an agent for a time and under conditions sufficient to modulate
 20 phosphorylation of said sphingosine kinase, which agent binds, links or otherwise associates with serine²²⁵, wherein inducing or otherwise agonising said phosphorylation up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates sphingosine kinase activity.

Yet another aspect of the present invention is directed to a method of modulating cellular
 25 activity, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-
 30 regulates said cellular activity.

A further aspect of the present invention is directed to a method of modulating cellular activity, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of human sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-
 5 regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

In another further aspect, the present invention is directed to a method of modulating human cellular activity, said method comprising contacting said cell with an effective
 10 amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of sphingosine kinase at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰ wherein inducing or otherwise agonising said phosphorylation up-regulates said human cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said human cellular activity.

15 Yet another aspect of the present invention is directed to a method for the treatment and/or prophylaxis of a condition in a mammal, which condition is characterised by aberrant, unwanted or otherwise inappropriate cellular activity, said method comprising administering to said mammal an effective amount of an agent for a time and under
 20 conditions sufficient to modulate phosphorylation of sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

25 In still another aspect, the present invention is directed to a method for the treatment and/or prophylaxis of a condition in a human, which condition is characterised by aberrant, unwanted or otherwise inappropriate cellular activity, said method comprising administering to said human an effective amount of an agent for a time and under
 30 conditions sufficient to modulate the phosphorylation of sphingosine kinase at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰ wherein inducing or otherwise agonising said

- 7 -

phosphorylation up-regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

Another aspect of the present invention contemplates the use of an agent, as hereinbefore
5 defined, in the manufacture of medicament for the treatment of a condition in a mammal,
which condition is characterised by aberrant, unwanted or otherwise inappropriate cellular
activity, wherein said agent modulates the phosphorylation of sphingosine kinase and
wherein inducing or otherwise agonising said phosphorylation up-regulates said cellular
activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said
10 cellular activity.

In yet another further aspect, the present invention contemplates a pharmaceutical
composition comprising the modulatory agent as hereinbefore defined together with one or
more pharmaceutically acceptable carriers and/or diluents. These agents are referred to as
15 the active ingredients.

Yet another aspect of the present invention relates to the agent as hereinbefore defined,
when used in the method of the present invention.

20 A further aspect of the present invention provides a method for detecting an agent capable
of modulating the phosphorylation of sphingosine kinase or its functional equivalent or
derivative thereof said method comprising contacting a cell or extract thereof containing
said sphingosine kinase and phosphorylation catalyst or its functional equivalent or
derivative with a putative agent and detecting an altered expression phenotype associated
25 with said phosphorylation.

Still another aspect of the present invention is directed to agents identified in accordance
with the screening method defined herein and to said agents for use in the methods of the
present invention. Said agent should be understood to extend to monoclonal antibodies
30 which bind to the phosphorylation sites of sphingosine kinase, and in particular, amino
acids S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ and T²⁵⁰, and most particularly S²²⁵.

Still a further aspect of the present invention is directed to sphingosine kinase variants comprising a mutation in a region of said sphingosine kinase which region comprises a phosphorylation site, wherein said variant exhibits ablated or reduced phosphorylation capacity relative to wild type sphingosine kinase or a derivative, homologue, analogue,
5 chemical equivalent or pneumatic of said sphingosine kinase variant.

In another aspect there is provided a human sphingosine kinase variant comprising an amino acid sequence with a single or multiple amino acid substitution and/or deletion of
10 amino acids S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ and/or T²⁵⁰ wherein said variant exhibits ablated or reduced phosphorylation capacity relative to wild-type sphingosine kinase or a derivative, homologue, analogue, chemical equivalent or mimetic of said sphingosine kinase variant.

In yet another aspect, the present invention extends to genetically modified animals, which
15 animals have been modified to express a sphingosine kinase variant as hereinbefore defined.

Single and three letter abbreviations used throughout the specification are defined in Table 1.

TABLE 1
Single and three letter amino acid abbreviations

5	Amino Acid	Three-letter	One-letter
		Abbreviation	Symbol
	Alanine	Ala	A
	Arginine	Arg	R
	Asparagine	Asn	N
10	Aspartic acid	Asp	D
	Cysteine	Cys	C
	Glutamine	Gln	Q
	Glutamic acid	Glu	E
	Glycine	Gly	G
15	Histidine	His	H
	Isoleucine	Ile	I
	Leucine	Leu	L
	Lysine	Lys	K
	Methionine	Met	M
20	Phenylalanine	Phe	F
	Proline	Pro	P
	Serine	Ser	S
	Threonine	The	T
	Tryptophan	Trp	W
25	Tyrosine	Tyr	Y
	Valine	Val	V
	Any residue	Xaa	X

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graphical representation of the phosphorylation of hSK1 paralleling its activation. TNF α stimulation of hSK1-transfected HEK293T cells resulted in an *in vivo* phosphorylation of hSK1 which was paralleled by an increase in sphingosine kinase activity in these cells.

Figure 2 is an image of the inactivation and lack of phosphorylation, in response to TNF α , of a mutant which is unable to bind TRAF2. Unlike wild-type hSK1, hSK^{TB2}, a hSK1 mutant defective in its ability to bind TRAF2 (Xia *et al.* 2002) is neither activated nor phosphorylated in response to TNF α treatment of HEK293T cells. In contrast, PMA treatment of these cells results in both phosphorylation and activation of wild-type hSK1 and hSK^{TB2}.

Figure 3 is a schematic representation of the prediction of phosphorylated amino acids in hSK1 using NetPhos. Circled residues represent those amino acids predicted to be phosphorylated in hSK1 (and its mouse and monkey homologues) by the NetPhos program.

Figure 4 is an image of the ablation of phosphorylation of hSK1 expressing an Ser²²⁵ \rightarrow Ala mutation. Metabolic labelling of HEK293T cells transfected with wild-type hSK1, hSK^{S148A}, hSK^{S181A}, hSK^{Y184A}, hSK^{S225A}, hSK^{T250A}, or empty vector and either untreated or treated with PMA showed that only the Ser²²⁵ \rightarrow Ala mutation ablated phosphorylation of hSK1.

25

Figure 5 is an image of the phosphorylation of hSK1, expressing an Ser²²⁵ \rightarrow Ala mutation, at adjacent sites. Metabolic labelling of HEK293T cells transfected with wild-type hSK1, hSK^{S220A}, hSK^{S225A}, or hSK^{T222A} and either untreated or treated with PMA again showed that only the Ser²²⁵ \rightarrow Ala mutation ablated phosphorylation of hSK1. This indicated that the Ser²²⁵ \rightarrow Ala mutation directly effects phosphorylation of hSK1 at Ser²²⁵ rather than indirectly effecting phosphorylation at adjacent sites (i.e. Ser²²⁰ and Thr²²²)

30

5

10

15

20

25

30

Figure 12 is a schematic representation of the targets for sphingosine kinase antagonists.

Figure 13 is a graphical representation of an ELISA of phospho-hSK antiserum. Crude open symbols) and affinity purified (closed symbols) antiserum from rabbits injected with a KLH-conjugated phosphopeptide designed around Ser²²⁵ of hSK1 was analysed by ELISA with the phosphopeptide (circles) and corresponding non-phosphorylated peptide (squares).

Figure 14 is an image illustrating that the phospho-hSK1 antiserum specifically reacts with phosphorylated hSK1 in Western blot analysis. Western blot analysis of hSK1 with the affinity purified phospho-hSK1 antiserum. Lane 1, recombinant hSK1; Lane 2, recombinant hSK1 phosphorylated *in vitro* by ERK2; Lane 3, recombinant hSK1 phosphorylated *in vitro* by ERK2 and subsequently dephosphorylated by alkaline phosphatase.

Figure 15 is an image illustrating that the phospho-hSK1 antiserum allows phosphorylation of *in vivo* hSK1 to be followed. HEK293T cells were transfected with wild-type hSK1 or hSK^{S225A} and treated with TNF α or PMA, harvested and analysed by Western blot with the phospho-hSK1 antiserum and anti-FLAG antibody.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is predicated, in part, on the determination that sphingosine kinase activation is mediated by a phosphorylation event and, further, the identification of the sphingosine kinase phosphorylation site itself. The inventors have still further determined that sphingosine kinase phosphorylation can be mediated by members of the proline-directed protein kinase family and, most preferably, ERK2. These determinations now permit the rational design of therapeutic and/or prophylactic methods for treating conditions, such as those characterised by aberrant or unwanted cellular activity and/or sphingosine kinase functional activity. Further, there is facilitated the identification and/or design of agents which specifically modulate sphingosine kinase phosphorylation.

Accordingly, one aspect of the present invention provides a method of modulating sphingosine kinase functional activity, said method comprising contacting said sphingosine kinase with an effective amount of an agent for a time and under conditions sufficient to modulate phosphorylation of said sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates sphingosine kinase activity.

Reference to "sphingosine kinase functional activity" should be understood as a reference to any one or more of the activities which sphingosine kinase can perform, for example, its activity as a key regulatory enzyme in the functioning of the sphingosine kinase signalling pathway. In this regard, it is thought that sphingosine kinase is central to the generation of sphingosine-1-phosphate during activation of this pathway. It should be understood that modulation of sphingosine kinase functional activity encompasses both up and down-regulation of any one or more of the functional activities attributable to sphingosine kinase, such as the induction or cessation of a given activity or a change to the level or degree of any given activity. Without limiting the invention to any one theory or mode of action, sphingosine kinase is thought to exhibit two levels of catalytic activity. At the first level, sphingosine kinase exhibits baseline catalytic activity. At the second level, sphingosine

- 14 -

kinase exhibiting baseline activity can be activated such that the Vmax of the enzyme is increased. In the context of the preferred embodiment of the present invention, the ablation or reduction of sphingosine kinase catalytic activity will be achieved where the baseline activity and/or the activation of sphingosine kinase beyond that of baseline
5 activity is ablated or reduced. Preferably, activation is ablated or reduced and even more preferably activation is ablated.

In a preferred embodiment, antagonising said phosphorylation prevents activation of sphingosine kinase and agonising said phosphorylation results in activation of said
10 sphingosine kinase.

Reference to "sphingosine kinase" should be understood to include reference to all forms of sphingosine kinase protein or derivatives, homologues, analogues, equivalents or mimetics thereof. In this regard, "sphingosine kinase" should be understood as being a
15 molecule which is, *inter alia*, involved in the generation of sphingosine-1-phosphate during activation of the sphingosine kinase signalling pathway. This includes, for example, all protein forms of sphingosine kinase or its functional derivatives, homologues, analogues, equivalents or mimetics thereof, including, for example, any isoforms which arise from alternative splicing of sphingosine kinase mRNA or allelic or polymorphic
20 variants of sphingosine kinase. Preferably, said sphingosine kinase is human sphingosine kinase.

Accordingly, the present invention provides a method of modulating human sphingosine kinase functional activity, said method comprising contacting said human sphingosine
25 kinase with an effective amount of an agent for a time and under conditions sufficient to modulate phosphorylation of said human sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said human sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said human sphingosine kinase activity.

30

Reference to "phosphorylation" should be understood as a reference to the addition of a phosphate group to the hydroxyl groups on proteins, in particular to the side chains of the amino acids serine, threonine or tyrosine. Without limiting the present invention to any one theory or mode of action, this process of phosphorylating proteins is normally
5 catalysed by a protein kinase (such molecules herein being referred to as "phosphorylation catalysts"), often a specific protein kinase, with ATP acting as the phosphate donor. The phosphorylation of proteins is generally found to regulate the activity of the subject protein.

10 As detailed hereinbefore, the phosphorylation of a protein is an event which occurs in the context of specific amino acid residues of a subject protein. In this regard, the inventors have identified S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ and T²⁵⁰ as being the amino acid residues which are relevant to phosphorylation of the sphingosine kinase molecule, in general. However, in terms of the human sphingosine kinase molecule, the inventors have still further identified
15 S²²⁵ as the primary physiological phosphorylation site of this molecule.

Accordingly, in a preferred embodiment, the present invention provides a method of modulating sphingosine kinase functional activity, said method comprising contacting said sphingosine kinase with an effective amount of an agent for a time and under conditions
20 sufficient to modulate the phosphorylation of said sphingosine kinase at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰, wherein inducing or otherwise agonising said phosphorylation up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said sphingosine kinase activity.

25 In a most preferred embodiment, said sphingosine kinase is human sphingosine kinase and said phosphorylation modulation occurs at the amino acid residue S²²⁵.

Reference to "modulating" either sphingosine kinase functional activity or a phosphorylation event should be understood as a reference to up-regulating or down-regulating the subject functional activity or phosphorylation event. Reference to up-regulating and down-regulating in this regard should be understood to include both
30

increasing or decreasing the level, degree or rate at which the functional activity or phosphorylation event occurs, in addition to including reference to inducing or ablating the subject functional activity or phosphorylation event. Accordingly, the agent which is utilised in accordance with the method of the present invention may be an agent which
5 induces the activity/event, agonises an activity or event which has already undergone onset, antagonises a pre-existing activity or event or entirely ablates such an activity or event.

Accordingly, reference to "inducing or otherwise agonising" phosphorylation should be
10 understood as a reference to:

- (i) inducing the phosphorylation of sphingosine kinase, for example, inducing the interaction of sphingosine kinase with a proline-directed protein kinase, such as ERK2, which effects sphingosine kinase phosphorylation; or
- 15 (ii) up-regulating, enhancing or otherwise agonising an existing phosphorylation event, for example, increasing the affinity of or otherwise stabilising the interaction of sphingosine kinase with a phosphorylating molecule.

20 Conversely, "inhibiting or otherwise antagonising" phosphorylation should be understood as a reference to:

- (i) preventing the interaction of sphingosine kinase with a phosphorylating molecule; or
- 25 (ii) antagonising an existing interaction between sphingosine kinase and a phosphorylating molecule such that the phosphorylation of sphingosine kinase is rendered ineffective or less effective.

It should be understood that modulation (either in the sense of up-regulation or down-
30 regulation) of the phosphorylation of sphingosine kinase may be partial or complete.

Partial modulation occurs where only some of the sphingosine kinase phosphorylation events which would normally occur in a given cell or on a given molecule are affected by the method of the present invention while complete modulation occurs where all sphingosine kinase phosphorylation events are modulated.

5

Modulation of the phosphorylation of sphingosine kinase may be achieved by any one of a number of techniques including, but not limited to:

- 10 (i) introducing into a cell a proteinaceous or non-proteinaceous agent which antagonises the interaction between sphingosine kinase and a phosphorylation catalyst;
- (ii) introducing into a cell a proteinaceous or non-proteinaceous agent which agonises the interaction between sphingosine kinase and a phosphorylation catalyst;
- 15 (iii) introducing into a cell a proteinaceous or non-proteinaceous agent which acts as a phosphorylation catalyst;
- (iv) introducing into a cell a nucleic acid molecule which encodes an agent which acts as a phosphorylation catalyst.
- 20

Preferably, said phosphorylation occurs at Ser²²⁵ of sphingosine kinase, and even more preferably, at Ser²²⁵ of human sphingosine kinase.

- 25 Reference to "agent" should be understood as a reference to any proteinaceous or non-proteinaceous molecule which modulates (i.e. up-regulates or down-regulates) the interaction of sphingosine kinase with a phosphorylation catalyst, for example, the molecules detailed in points (i) – (ii), above. "Agent" should also be understood to extend to molecules which, of themselves, function as a phosphorylation catalyst, for example, the
- 30 molecules detailed in (iii) - (iv), above. The subject agent may be linked, bound or

otherwise associated with any proteinaceous or non-proteinaceous molecule. For example, it may be associated with a molecule which permits targeting to a localised region.

- Said proteinaceous molecule may be derived from natural, recombinant or synthetic
- 5 sources including fusion proteins or following, for example, natural product screening.
- Said non-proteinaceous molecule may be derived from natural sources, such as for example natural product screening or may be chemically synthesised. For example, the present invention contemplates chemical analogues of sphingosine kinase or
- 10 phosphorylation catalyst, such as a kinase molecule (for example, a proline-directed protein kinase such as ERK2), capable of acting as agonists or antagonists of sphingosine kinase phosphorylation. Chemical agonists may not necessarily be derived from sphingosine kinase or the phosphorylation catalyst but may share certain conformational similarities. Alternatively, chemical agonists may be specifically designed to mimic certain physiochemical properties of sphingosine kinase or phosphorylation catalyst.
- 15 Antagonists may be any compound capable of blocking, inhibiting or otherwise preventing sphingosine kinase phosphorylation. Antagonists include monoclonal antibodies specific for sphingosine kinase or the phosphorylation catalyst, or parts of said sphingosine kinase, and antisense nucleic acids which prevent transcription or translation of genes or mRNA in the subject cells. Modulation of expression may also be achieved utilising antigens, RNA,
- 20 ribosomes, DNazymes, RNA aptamers, antibodies or molecules suitable for use in co-suppression. Screening methods suitable for use in identifying such molecules are described in more detail hereinafter.

- Said proteinaceous or non-proteinaceous molecule may act either directly or indirectly to
- 25 modulate the interaction of sphingosine kinase with a phosphorylating molecule (such as a phosphorylating kinase). Said molecule acts directly if it associates with the sphingosine kinase or the phosphorylation catalyst. Said molecule acts indirectly if it associates with a molecule other than the sphingosine kinase or phosphorylation catalyst, which other molecule either directly or indirectly modulates the interaction of sphingosine kinase with
- 30 the phosphorylation catalyst. Accordingly, the method of the present invention

encompasses regulation of sphingosine kinase phosphorylation via the induction of a cascade of regulatory steps.

It should be understood that reference to "phosphorylation catalyst" is intended to encompass both molecules which catalyse the transfer of phosphate from a molecule such as ATP to sphingosine kinase. However, it should also be understood to encompass reference to molecules which otherwise achieve the phosphorylation of sphingosine kinase, such as molecules which are able to transfer phosphate groups from molecules other than ATP or which can directly transfer a phosphate group from the phosphorylation catalyst itself to the sphingosine kinase molecule.

"Derivatives" include fragments, parts, portions, mutants, variants and mimetics from natural, synthetic or recombinant sources including fusion proteins. Parts or fragments include, for example, active regions of sphingosine kinase or the phosphorylation catalyst. Derivatives may be derived from insertion, deletion or substitution of amino acids. Amino acid insertional derivatives include amino and/or carboxylic terminal fusions as well as intrasequence insertions of single or multiple amino acids. Insertional amino acid sequence variants are those in which one or more amino acid residues are introduced into a predetermined site in the protein although random insertion is also possible with suitable screening of the resulting product. Deletional variants are characterized by the removal of one or more amino acids from the sequence. Substitutional amino acid variants are those in which at least one residue in the sequence has been removed and a different residue inserted in its place. An example of substitutional amino acid variants are conservative amino acid substitutions. Conservative amino acid substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine and leucine; aspartic acid and glutamic acid; asparagine and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine. Additions to amino acid sequences include fusions with other peptides, polypeptides or proteins.

Reference to "homologues" should be understood as a reference to nucleic acid molecules or proteins derived from alternative species.

Equivalents of nucleic acid or protein molecules should be understood as molecules exhibiting any one or more of the functional activities of these molecules and may be derived from any source such as being chemically synthesized or identified via screening
5 processes such as natural product screening.

The derivatives include fragments having particular epitopes or parts of the entire protein fused to peptides, polypeptides or other proteinaceous or non-proteinaceous molecules.

10. Analogues contemplated herein include, but are not limited to, modification to side chains, incorporating of unnatural amino acids and/or their derivatives during peptide, polypeptide or protein synthesis and the use of crosslinkers and other methods which impose conformational constraints on the proteinaceous molecules or their analogues.

15 Derivatives of nucleic acid sequences may similarly be derived from single or multiple nucleotide substitutions, deletions and/or additions including fusion with other nucleic acid molecules. The derivatives of the nucleic acid molecules of the present invention include oligonucleotides, PCR primers, antisense molecules, molecules suitable for use in cosuppression and fusion of nucleic acid molecules. Derivatives of nucleic acid sequences
20 also include degenerate variants.

Examples of side chain modifications contemplated by the present invention include modifications of amino groups such as by reductive alkylation by reaction with an aldehyde followed by reduction with NaBH_4 ; amidination with methylacetimidate;
25 acylation with acetic anhydride; carbamoylation of amino groups with cyanate; trinitrobenzylation of amino groups with 2, 4, 6-trinitrobenzene sulphonic acid (TNBS); acylation of amino groups with succinic anhydride and tetrahydrophthalic anhydride; and pyridoxylation of lysine with pyridoxal-5-phosphate followed by reduction with NaBH_4 .

30 The guanidine group of arginine residues may be modified by the formation of heterocyclic condensation products with reagents such as 2,3-butanedione, phenylglyoxal and glyoxal.

The carboxyl group may be modified by carbodiimide activation *via* O-acylisourea formation followed by subsequent derivitisation, for example, to a corresponding amide.

5 Sulphydryl groups may be modified by methods such as carboxymethylation with iodoacetic acid or iodoacetamide; performic acid oxidation to cysteic acid; formation of a mixed disulphides with other thiol compounds; reaction with maleimide, maleic anhydride or other substituted maleimide; formation of mercurial derivatives using 4-chloromercuribenzoate, 4-chloromercuriphenylsulphonic acid, phenylmercury chloride, 2-chloromercuri-4-nitrophenol and other mercurials; carbamoylation with cyanate at alkaline
10 pH.

Tryptophan residues may be modified by, for example, oxidation with N-bromosuccinimide or alkylation of the indole ring with 2-hydroxy-5-nitrobenzyl bromide or sulphenyl halides. Tyrosine residues on the other hand, may be altered by nitration with
15 tetranitromethane to form a 3-nitrotyrosine derivative.

Modification of the imidazole ring of a histidine residue may be accomplished by alkylation with iodoacetic acid derivatives or N-carboethoxylation with diethylpyrocarbonate.

20 Examples of incorporating unnatural amino acids and derivatives during protein synthesis include, but are not limited to, use of norleucine, 4-amino butyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, ornithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid, 2-thienyl
25 alanine and/or D-isomers of amino acids. A list of unnatural amino acids contemplated herein is shown in Table 2.

TABLE 2

5	Non-conventional amino acid	Code	Non-conventional amino acid	Code
	α -aminobutyric acid	Abu	L-N-methylalanine	Nmala
	α -amino- α -methylbutyrate	Mgab	L-N-methylarginine	Nmarg
10	aminocyclopropane- carboxylate	Cpro	L-N-methylasparagine	Nmasn
	aminoisobutyric acid	Aib	L-N-methylaspartic acid	Nmasp
	aminonorbornyl- carboxylate	Norb	L-N-methylcysteine	Nmcys
			L-N-methylglutamine	Nmgln
			L-N-methylglutamic acid	Nmglu
15	cyclohexylalanine	Chexa	L-N-methylhistidine	Nmhis
	cyclopentylalanine	Cpen	L-N-methylisoleucine	Nmile
	D-alanine	Dal	L-N-methylleucine	Nmleu
	D-arginine	Darg	L-N-methyllysine	Nmlys
	D-aspartic acid	Dasp	L-N-methylmethionine	Nmmet
20	D-cysteine	Dcys	L-N-methylnorleucine	Nmnle
	D-glutamine	Dgln	L-N-methylnorvaline	Nmnva
	D-glutamic acid	Dglu	L-N-methylornithine	Nmorn
	D-histidine	Dhis	L-N-methylphenylalanine	Nmphe
	D-isoleucine	Dile	L-N-methylproline	Nmpro
25	D-leucine	Dleu	L-N-methylserine	Nmser
	D-lysine	Dlys	L-N-methylthreonine	Nmthr
	D-methionine	Dmet	L-N-methyltryptophan	Nmtrp
	D-ornithine	Dorn	L-N-methyltyrosine	Nmtyr
	D-phenylalanine	Dphe	L-N-methylvaline	Nmval
30	D-proline	Dpro	L-N-methylethylglycine	Nmetg
	D-serine	Dser	L-N-methyl-t-butylglycine	Nmtbug
	D-threonine	Dthr	L-norleucine	Nle

- 23 -

	D-tryptophan	Dtrp	L-norvaline	Nva
	D-tyrosine	Dtyr	α -methyl-aminoisobutyrate	Maib
	D-valine	Dval	α -methyl- -aminobutyrate	Mgab
	D- α -methylalanine	Dmala	α -methylcyclohexylalanine	Mchexa
5	D- α -methylarginine	Dmarg	α -methylcyclopentylalanine	Mcpen
	D- α -methylasparagine	Dmasn	α -methyl- α -naphthylalanine	Manap
	D- α -methylaspartate	Dmasp	α -methylpenicillamine	Mpen
	D- α -methylcysteine	Dmcys	N-(4-aminobutyl)glycine	Nglu
	D- α -methylglutamine	Dmgln	N-(2-aminoethyl)glycine	Naeg
10	D- α -methylhistidine	Dmhis	N-(3-aminopropyl)glycine	Norn
	D- α -methylisoleucine	Dmile	N-amino- α -methylbutyrate	Nmaabu
	D- α -methylleucine	Dmleu	α -naphthylalanine	Anap
	D- α -methyllysine	Dmlys	N-benzylglycine	Nphe
	D- α -methylmethionine	Dmmet	N-(2-carbamylethyl)glycine	Ngln
15	D- α -methylornithine	Dmorn	N-(carbamylmethyl)glycine	Nasn
	D- α -methylphenylalanine	Dmphe	N-(2-carboxyethyl)glycine	Nglu
	D- α -methylproline	Dmpro	N-(carboxymethyl)glycine	Nasp
	D- α -methylserine	Dmser	N-cyclobutylglycine	Ncbut
	D- α -methylthreonine	Dmthr	N-cycloheptylglycine	Nchep
20	D- α -methyltryptophan	Dmtrp	N-cyclohexylglycine	Nchex
	D- α -methyltyrosine	Dmtty	N-cyclodecylglycine	Ncdec
	D- α -methylvaline	Dmval	N-cylcododecylglycine	Ncdod
	D-N-methylalanine	Dnmala	N-cyclooctylglycine	Ncoct
	D-N-methylarginine	Dnmarg	N-cyclopropylglycine	Ncpro
25	D-N-methylasparagine	Dnmasn	N-cycloundecylglycine	Ncund
	D-N-methylaspartate	Dnmasp	N-(2,2-diphenylethyl)glycine	Nbhm
	D-N-methylcysteine	Dnmcys	N-(3,3-diphenylpropyl)glycine	Nbhe
	D-N-methylglutamine	Dnmgln	N-(3-guanidinopropyl)glycine	Narg
	D-N-methylglutamate	Dnmglu	N-(1-hydroxyethyl)glycine	Nthr
30	D-N-methylhistidine	Dnmhis	N-(hydroxyethyl)glycine	Nser
	D-N-methylisoleucine	Dnmile	N-(imidazolylethyl)glycine	Nhis
	D-N-methylleucine	Dnmleu	N-(3-indolylyethyl)glycine	Nhtrp
	D-N-methyllysine	Dnmlys	N-methyl- γ -aminobutyrate	Nmgabu

	N-methylcyclohexylalanine	Nmchexa	D-N-methylmethionine	Dnmmt
	D-N-methylornithine	Dnmorn	N-methylcyclopentylalanine	Nmcpn
	N-methylglycine	Nala	D-N-methylphenylalanine	Dnmph
	N-methylaminoisobutyrate	Nmaib	D-N-methylproline	Dnmpro
5	N-(1-methylpropyl)glycine	Nile	D-N-methylserine	Dnmser
	N-(2-methylpropyl)glycine	Nleu	D-N-methylthreonine	Dnmthr
	D-N-methyltryptophan	Dnmtrp	N-(1-methylethyl)glycine	Nval
	D-N-methyltyrosine	Dnmtyr	N-methyl- <i>n</i> -naphthylalanine	Nmanap
	D-N-methylvaline	Dnmval	N-methylpenicillamine	Nmpen
10	γ -aminobutyric acid	Gabu	N-(<i>p</i> -hydroxyphenyl)glycine	Nhtyr
	L- <i>t</i> -butylglycine	Tbug	N-(thiomethyl)glycine	Ncys
	L-ethylglycine	Etg	penicillamine	Pen
	L-homophenylalanine	Hphe	L- α -methylalanine	Mala
	L- α -methylarginine	Marg	L- α -methylnorleucine	Masn
15	L- α -methyloxyaspartate	Masp	L- α -methyl- <i>t</i> -butylglycine	Mtbug
	L- α -methylcysteine	Mcys	L-methylethylglycine	Metg
	L- α -methylglutamine	Mgin	L- α -methylglutamate	Mglu
	L- α -methylhistidine	Mhis	L- α -methylhomophenylalanine	Mhphe
	L- α -methylisoleucine	Mile	N-(2-methylthioethyl)glycine	Nmet
20	L- α -methylleucine	Mleu	L- α -methyllysine	Mlys
	L- α -methylmethionine	Mmet	L- α -methylnorleucine	Mnle
	L- α -methylnorvaline	Mnva	L- α -methylornithine	Morn
	L- α -methylphenylalanine	Mphe	L- α -methylproline	Mpro
	L- α -methylserine	Mser	L- α -methylthreonine	Mthr
25	L- α -methyltryptophan	Mtrp	L- α -methyltyrosine	Mtyr
	L- α -methylvaline	Mval	L-N-methylhomophenylalanine	Nmhph
	N-(N-(2,2-diphenylethyl)carbamylmethyl)glycine	Nnbhm	N-(N-(3,3-diphenylpropyl)carbamylmethyl)glycine	Nnbhe
30	1-carboxy-1-(2,2-diphenyl)ethylamino)cyclopropane			

Crosslinkers can be used, for example, to stabilise 3D conformations, using homo-bifunctional crosslinkers such as the bifunctional imido esters having $(CH_2)_n$ spacer groups with $n=1$ to $n=6$, glutaraldehyde, N-hydroxysuccinimide esters and hetero-bifunctional reagents which usually contain an amino-reactive moiety such as N-hydroxysuccinimide
5 and another group specific-reactive moiety.

Without limiting the present invention to any one theory or mode of action, the inventors have utilised alanine mutagenesis to determine that serine ²²⁵ is the primary physiologic phosphorylation site on human sphingosine kinase 1. Further, it has been determined that
10 the kinases of the proline-directed protein kinase family are able to phosphorylate human sphingosine kinase. Specifically, ERK1, ERK2 and CDK2 phosphorylate human sphingosine kinase 1 to varying degrees of efficiency. ERK2, however, has been demonstrated to show the greatest efficiency in human sphingosine kinase phosphorylation.

15 Accordingly, in a most preferred embodiment the present invention is directed to a method of modulating sphingosine kinase functional activity, said method comprising contacting said sphingosine kinase with an effective amount of an agent for a time and under conditions sufficient to modulate proline-directed protein kinase catalysed sphingosine
20 kinase phosphorylation wherein inducing or otherwise agonising said phosphorylation up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said phosphorylation down-regulates sphingosine kinase activity.

Most preferably, said proline directed protein kinase is ERK1, ERK2 or CDK2 and, even
25 more particularly, ERK2.

Most preferably, said phosphorylation is modulated at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰ and most particularly at S²²⁵.

30 Reference to proline-directed protein kinase molecules, such as ERK1, ERK2 and CDK2, and any other phosphorylation catalyst should be understood to encompass reference to

derivatives, homologues, analogues, chemical equivalents and mimetics of these molecules.

In a related aspect the present invention provides a method of modulating sphingosine
5 kinase functional activity, said method comprising contacting said sphingosine kinase with
an effective amount of an agent for a time and under conditions sufficient to modulate
phosphorylation of said sphingosine kinase, which agent binds, links or otherwise
associates with serine ²²⁵, wherein inducing or otherwise agonising said phosphorylation
up-regulates said sphingosine kinase activity and inhibiting or otherwise antagonising said
10 phosphorylation down-regulates sphingosine kinase activity.

In a most preferred embodiment, said modulation of sphingosine kinase activity is down-
regulation.

15 Since sphingosine kinase is a molecule which is central to the functioning of an
intracellular signalling pathway, the method of the present invention provides a means of
modulating cellular activity which is regulated or controlled by sphingosine kinase
signalling. For example, the sphingosine kinase signalling pathway is known to regulate
cellular activities, such as those which lead to inflammation, apoptosis and cell
20 proliferation (for example, up-regulation of the production of inflammatory mediators such
as cytokines, chemokines, eNOS and up-regulation of adhesion molecule expression).
Said up-regulation may be induced by a number of stimuli including, for example,
inflammatory cytokines such as tumour necrosis factor α and interleukin 1, endotoxin,
oxidised or modified lipids, radiation or tissue injury. In this regard, reference to
25 "modulating cellular activity" is a reference to up-regulating, down-regulating or otherwise
altering any one or more of the activities which a cell is capable of performing pursuant to
sphingosine kinase signalling such as, but not limited, one or more of chemokine
production, cytokine production, nitric oxide synthesis, adhesion molecule expression and
production of other inflammatory modulators. Although the preferred method is to down-
30 regulate sphingosine kinase activity, thereby down-regulating unwanted cellular activity,

the present invention should nevertheless be understood to encompass up-regulating of cellular activity, which may be desirable in certain circumstances.

Accordingly, yet another aspect of the present invention is directed to a method of modulating cellular activity, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

Preferably, the present invention is directed to a method of modulating cellular activity, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of human sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

In a most preferred embodiment the present invention is directed to a method of modulating human cellular activity, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of sphingosine kinase at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰ wherein inducing or otherwise agonising said phosphorylation up-regulates said human cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said human cellular activity.

Most preferably, said phosphorylation modulation occurs at amino acid residue S²²⁵ and said modulation is down-regulation.

A further aspect of the present invention relates to the use of the invention in relation to the treatment and/or prophylaxis of disease conditions. Without limiting the present invention to any one theory or mode of action, the broad range of cellular functional activities which

are regulated via the sphingosine kinase signalling pathway renders the regulation of sphingosine kinase functioning an integral component of every aspect of both healthy and disease state physiological processes. Accordingly, the method of the present invention provides a valuable tool for modulating aberrant or otherwise unwanted cellular functional
5 activity which is regulated via the sphingosine kinase signalling pathway.

Accordingly, yet another aspect of the present invention is directed to a method for the treatment and/or prophylaxis of a condition in a mammal, which condition is characterised by aberrant, unwanted or otherwise inappropriate cellular activity, said method comprising
10 administering to said mammal an effective amount of an agent for a time and under conditions sufficient to modulate phosphorylation of sphingosine kinase wherein inducing or otherwise agonising said phosphorylation up-regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

15 Preferably, said phosphorylation event is phosphorylation of sphingosine kinase at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰.

In a most preferred embodiment, the present invention is directed to a method for the treatment and/or prophylaxis of a condition in a human, which condition is characterised by aberrant, unwanted or otherwise inappropriate cellular activity, said method comprising
20 administering to said human an effective amount of an agent for a time and under conditions sufficient to modulate the phosphorylation of sphingosine kinase at one or more of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰ wherein inducing or otherwise agonising said
25 phosphorylation up-regulates said cellular activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

Most preferably, said phosphorylation event is phosphorylation of sphingosine kinase at S²²⁵ and still more preferably, said modulation is down-regulation.

Reference to "aberrant, unwanted or otherwise inappropriate" cellular activity should be understood to be understood as a reference to overactive cellular activity, to physiological normal cellular activity which is inappropriate in that it is unwanted or to insufficient cellular activity. This definition applies in an analogous manner in relation to "aberrant, unwanted or otherwise, inappropriate" sphingosine kinase activation. For example, TNF production during tumour cell growth has been shown to support cellular proliferation and to provide anti-apoptotic characteristics to the neo plastic cells. Accordingly, to the extent that a cell is neoplastic, it is desirable that the promotion of cellular proliferation and anti-apoptotic characteristics be down-regulated. Similarly, diseases which are characterised by inflammation, such as rheumatoid arthritis, are known to involve cellular activation by cytokines such as TNF, leading to the synthesis and secretion of inflammatory mediators. In such situations, it is also desirable to down-regulate such activity. In other situations, it may be desirable to agonise or otherwise induce sphingosine kinase phosphorylation in order to stimulate cellular proliferation.

In accordance with the previous aspects of the present invention, the subject agent is preferably PD98059 or derivative, analogue, chemical equivalent or mimetic thereof.

The term "mammal" as used herein includes humans, primates, livestock animals (eg. sheep, pigs, cattle, horses, donkeys), laboratory test animals (eg. mice, rabbits, rats, guinea pigs), companion animals (eg. dogs, cats) and captive wild animals (eg. foxes, kangaroos, deer). Preferably, the mammal is human or a laboratory test animal. Even more preferably, the mammal is a human.

An "effective amount" means an amount necessary at least partly to attain the desired response, or to delay the onset or inhibit progression or halt altogether, the onset or progression of a particular condition being treated. The amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated, the degree of protection desired, the formulation of the composition, the assessment of the medical situation, and other relevant factors. It is

expected that the amount will fall in a relatively broad range that can be determined through routine trials.

Reference herein to "treatment" and "prophylaxis" is to be considered in its broadest context. The term "treatment" does not necessarily imply that a subject is treated until total recovery. Similarly, "prophylaxis" does not necessarily mean that the subject will not eventually contract a disease condition. Accordingly, treatment and prophylaxis include amelioration of the symptoms of a particular condition or preventing or otherwise reducing the risk of developing a particular condition. The term "prophylaxis" may be considered as reducing the severity or onset of a particular condition. "Treatment" may also reduce the severity of an existing condition.

The present invention further contemplates a combination of therapies, such as the administration of the agent together with subjection of the mammal to other agents, drugs or treatments which may be useful in relation to the treatment of the subject condition such as cytotoxic agents or radiotherapy in the treatment of cancer.

Administration of the modulatory agent, in the form of a pharmaceutical composition, may be performed by any convenient means. The modulatory agent of the pharmaceutical composition is contemplated to exhibit therapeutic activity when administered in an amount which depends on the particular case. The variation depends, for example, on the human or animal and the modulatory agent chosen. A broad range of doses may be applicable. Considering a patient, for example, from about 0.1 mg to about 1 mg of modulatory agent may be administered per kilogram of body weight per day. Dosage regimes may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily, weekly, monthly or other suitable time intervals or the dose may be proportionally reduced as indicated by the exigencies of the situation.

The modulatory agent may be administered in a convenient manner such as by the oral, intravenous (where water soluble), intraperitoneal, intramuscular, subcutaneous,

intradermal or suppository routes or implanting (e.g. using slow release molecules). The modulatory agent may be administered in the form of pharmaceutically acceptable nontoxic salts, such as acid addition salts or metal complexes, e.g. with zinc, iron or the like (which are considered as salts for purposes of this application). Illustrative of such acid addition salts are hydrochloride, hydrobromide, sulphate, phosphate, maleate, acetate, citrate, benzoate, succinate, malate, ascorbate, tartrate and the like. If the active ingredient is to be administered in tablet form, the tablet may contain a binder such as tragacanth, corn starch or gelatin; a disintegrating agent, such as alginic acid; and a lubricant, such as magnesium stearate.

Routes of administration include, but are not limited to, respiratorally, intratracheally, nasopharyngeally, intravenously, intraperitoneally, subcutaneously, intracranially, intradermally, intramuscularly, intraocularly, intrathecally, intracerebrally, intranasally, infusion, orally, rectally, *via* IV drip patch and implant.

In accordance with these methods, the agent defined in accordance with the present invention may be coadministered with one or more other compounds or molecules. By "coadministered" is meant simultaneous administration in the same formulation or in two different formulations via the same or different routes or sequential administration by the same or different routes. For example, the subject agent may be administered together with an agonistic agent in order to enhance its effects. By "sequential" administration is meant a time difference of from seconds, minutes, hours or days between the administration of the two types of molecules. These molecules may be administered in any order.

Another aspect of the present invention contemplates the use of an agent, as hereinbefore defined, in the manufacture of medicament for the treatment of a condition in a mammal, which condition is characterised by aberrant, unwanted or otherwise inappropriate cellular activity, wherein said agent modulates the phosphorylation of sphingosine kinase and wherein inducing or otherwise agonising said phosphorylation up-regulates said cellular

activity and inhibiting or otherwise antagonising said phosphorylation down-regulates said cellular activity.

Preferably, said phosphorylation is phosphorylation of sphingosine kinase at one or more
5 of S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ or T²⁵⁰.

Even more preferably, said mammal is a human, said phosphorylation is phosphorylation of sphingosine kinase at S²²⁵ and said modulation is down-regulation.

10 As detailed hereinbefore, and without limiting the present invention to any one theory or mode of action, it is thought that phosphorylation of sphingosine kinase activates sphingosine kinase exhibiting baseline activity such that the V_{max} of the enzyme is increased. Accordingly, screening individuals for the presence, absence or degree of sphingosine kinase phosphorylation provides a means for diagnosing, monitoring or
15 screening patients for conditions characterised by aberrantly activated sphingosine kinase activity and/or cellular activity. The present invention should therefore be understood to extend to such diagnostic assays.

In yet another further aspect, the present invention contemplates a pharmaceutical
20 composition comprising the modulatory agent as hereinbefore defined together with one or more pharmaceutically acceptable carriers and/or diluents. These agents are referred to as the active ingredients.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions
25 (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion or may be in the form of a cream or other form suitable for topical application. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion
30 medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof, and

vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The preventions of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

10

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilisation. Generally, dispersions are prepared by incorporating the various sterilised active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

20

When the active ingredients are suitably protected they may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsule, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 1% by weight of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 5 to about 80% of the weight of the unit. The amount of active compound in such therapeutically useful compositions in such that a suitable dosage will be obtained. Preferred compositions or preparations according to the

30

- 34 -

present invention are prepared so that an oral dosage unit form contains between about 0.1 µg and 2000 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the components as listed hereafter: a binder such as gum, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound(s) may be incorporated into sustained-release preparations and formulations.

The pharmaceutical composition may also comprise genetic molecules such as a vector capable of transfecting target cells where the vector carries a nucleic acid molecule encoding a modulatory agent. The vector may, for example, be a viral vector.

Yet another aspect of the present invention relates to the agent as hereinbefore defined, when used in the method of the present invention.

25

The present invention should also be understood to encompass a method for screening for agents which modulate the phosphorylation of sphingosine kinase, particularly at Ser²²⁵, such as agents which agonise or antagonise the interaction of sphingosine kinase with a phosphorylation catalyst (such as ERK2) or which themselves phosphorylate sphingosine kinase, for example by functioning as a phosphorylation catalyst.

30

5

15

25

30

Accordingly, another aspect of the present invention provides a method for detecting an agent capable of modulating the phosphorylation of sphingosine kinase or its functional equivalent or derivative thereof said method comprising contacting a cell or extract thereof containing said sphingosine kinase and phosphorylation catalyst or its functional
5 equivalent or derivative with a putative agent and detecting an altered expression phenotype associated with said phosphorylation.

Reference to "sphingosine kinase" and "phosphorylation catalyst" should be understood as a reference to either the sphingosine kinase or phosphorylation catalyst expression product
10 or to a portion or fragment of the sphingosine kinase or phosphorylation catalyst, such as the Ser²²⁵ sphingosine kinase phosphorylation site. In this regard, the sphingosine kinase or phosphorylation catalyst expression product is expressed in a cell. The cell may be a host cell which has been transfected with the sphingosine kinase or phosphorylation catalyst nucleic acid molecule or it may be a cell which naturally contains the sphingosine
15 kinase gene. Reference to "extract thereof" should be understood as a reference to a cell free transcription system.

Reference to detecting an "altered expression phenotype associated with said phosphorylation" should be understood as the detection of cellular changes associated with
20 modulation of the phosphorylation of sphingosine kinase. These may be detectable, for example, as intracellular changes or changes observable extracellularly. For example, this includes, but is not limited to, detecting changes in downstream product levels or activities (e.g. NF-κB).

25 Still another aspect of the present invention is directed to agents identified in accordance with the screening method defined herein and to said agents for use in the methods of the present invention. Said agent should be understood to extend to monoclonal antibodies which bind to the phosphorylation sites of sphingosine kinase, and in particular, amino acids S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ and T²⁵⁰, and most particularly S²²⁵.

Still a further aspect of the present invention is directed to sphingosine kinase variants comprising a mutation in a region of said sphingosine kinase which region comprises a phosphorylation site, wherein said variant exhibits ablated or reduced phosphorylation capacity relative to wild type sphingosine kinase or a derivative, homologue, analogue,
5 chemical equivalent or pneumatic of said sphingosine kinase variant.

The present invention also extends to variants which exhibit enhanced or up-regulated activity due to the nature of the mutation of an existing phosphorylation site or the incorporation of additional phosphorylation sites.

10

Reference to "mutation" should be understood as a reference to any change, alteration or other modification, whether occurring naturally or non-naturally, which modulates the capacity of said sphingosine kinase to undergo phosphorylation. Said modulation may be up-regulation or down-regulation. Although the present invention is preferably directed to
15 variants which exhibit ablated activation capacity, it should be understood that the present invention extends to the generation of variants which exhibit additional phosphorylation sites or an improved level of phosphorylation capacity of existing sites.

The change, alteration or other modification may take any form including, but not limited
20 to, a structural modification (such an alteration in the secondary, tertiary or quaternary structure of the sphingosine kinase molecule), a molecular modification (such as an addition, substitution or deletion of one or more amino acids from the sphingosine kinase protein) or a chemical modification. The subject modification should also be understood to extend to the fusion, linking or binding of a proteinaceous or non-proteinaceous
25 molecule to the sphingosine kinase protein or to the nucleic acid molecule encoding a sphingosine kinase protein. It should also be understood that although it is necessary that the subject mutation is expressed by the sphingosine kinase expression product, the creation of the mutation may be achieved by any suitable means including either mutating a wild-type sphingosine kinase protein, synthesising a sphingosine kinase variant or
30 modifying a nucleic acid molecule encoding a wild-type sphingosine kinase protein such that the expression product of said mutated nucleic acid molecule is a sphingosine kinase

protein variant. Preferably, said mutation is a single or multiple amino acid sequence substitution, addition and/or deletion.

5 In accordance with this preferred embodiment there is provided a human sphingosine kinase variant comprising an amino acid sequence with a single or multiple amino acid substitution and/or deletion of amino acids S¹⁴⁸, S¹⁸¹, Y¹⁸⁴, S²²⁵ and/or T²⁵⁰ wherein said variant exhibits ablated or reduced phosphorylation capacity relative to wild-type sphingosine kinase or a derivative, homologue, analogue, chemical equivalent or mimetic of said sphingosine kinase variant.

10 Preferably, amino acid is amino acid S²²⁵ and even more preferably said mutation is a Ser²²⁵ Ala substitution.

15 In terms of the present invention, reference to "wild-type" sphingosine kinase is a reference to the forms of sphingosine kinase expressed by most individuals in a given population. There may be greater than one wild-type form of sphingosine kinase (for example due to allelic or isoform variation) and the level of phosphorylation exhibited by said wild-type sphingosine kinase molecules may fall within a range of levels. However, it should be understood that "wild-type" does not include reference to a naturally occurring form of sphingosine kinase which cannot be phosphorylated. Such a variant form of sphingosine kinase may, in fact, constitute a naturally occurring mutant form of sphingosine kinase within the context of the present invention.

25 In yet another aspect, the present invention extends to genetically modified animals, which animals have been modified to express a sphingosine kinase variant as hereinbefore defined.

The present invention is further described by reference to the following non-limiting description.

EXAMPLE 1

Methods

Generation of phospho-hSK1-specific polyclonal antibodies

5 Polyclonal antibodies were raised in rabbits against a phosphopeptide (CGSKTPApSPVVVQQ) designed from the hSK1 sequence around Ser²²⁵. Prior to injection into rabbits, the phosphopeptide was conjugated to maleimide activated keyhole limpet hemocyanin (Pierce) via its *N*-terminal cysteine. Antibodies active against the non-
10 phosphorylated peptide were removed from the antiserum using SulfoLink™ beads (Pierce) to which the non-phosphorylated peptide (CGSKTPASPVVVQQ <400>2) was conjugated using the manufactures instructions.

Construction and expression of hSK mutants

15 Wild-type human SK1 (hSKWT) cDNA (Pitson *et al.* 2000) (Genbank accession number AF200328) was FLAG epitope tagged at the 3' end and subcloned into pALTER (Promega Inc., Madison, WI) site directed mutagenesis vector, as previously described (Pitson *et al.* 2000). Single-stranded DNA was prepared and used as template for oligonucleotide
20 directed mutagenesis as detailed in the manufacturer's protocol. The mutagenic oligonucleotides shown in Table 1 were designed to generate the required mutants which were subsequently sequenced to verify incorporation of the desired modification. The mutant cDNA were then subcloned into pcDNA3 (Invitrogen Corp., San Diego, CA) for transient transfection into HEK293T cells of pGEX-4T2 for expression as glutathione-S-
25 transferase (GST)-fusion proteins in *E. coli*. hSK^{TB2} was generated as previously described (Xia *et al.* 2002).

Cell Culture and Transfection

30 Human embryonic kidney cells (HEK293T, ATCC CRL-1573) were cultured on Dulbecco's modified Eagle's medium (CSL Biosciences, Parkville, Australia) containing

10% fetal calf serum, 2 mM glutamine, 0.2% (w/v) sodium bicarbonate, penicillin (1.2 mg/ml), and gentamycin (1.6 mg/ml). Transfections were performed using the calcium phosphate precipitation method. Cells were harvested 24 h after transfection and lysed by sonication (2 watts for 30 s at 4 °C) in lysis buffer A containing 50 mM Tris/HCl (pH 7.4),
5 10% (w/v) glycerol, 0.05% (w/v) Triton X-100, 150 mM NaCl, 1 mM dithiothreitol, 2 mM Na₃VO₄, 10 mM NaF, 1 mM EDTA and protease inhibitors (Complete™; Boehringer Mannheim). Activation of hSK1 was assessed by treatment of cells with either phorbol 12-myristate 13-acetate (PMA; Sigma) for 30 min, or tumor necrosis factor- α (TNF α ; R&D Systems Inc., Minneapolis, MN) for 10 min. Protein concentrations in cell
10 homogenates were determined with Coomassie Brilliant Blue (Sigma) reagent using BSA as standard.

Generation of recombinant proteins in E. coli

15 Recombinant proteins were generated of in *E. coli* and purified as previously described (Pitson *et al.* 2000).

Metabolic labelling of cells with ³²P

20 HEK293T cells were cultured and transfected as described above. The cultures, in 10 cm dishes, were incubated for 30 min with phosphate-free, serum-free Dulbecco's modified Eagle's medium (DMEM). This medium was then replaced with fresh phosphate-free, serum-free DMEM containing 500 μ Ci [³²P]orthophosphate and the cultures incubated for 3 hr. Cultures were then stimulated with 1 ng/ml TNF or 100 ng/ml PMA for 30 min. Cells
25 were then washed 3 times with cold PBS and scraped into 1.2 ml of lysis buffer B containing 50 mM Tris/HCl (pH 7.4), 1 % (w/v) Triton X-100, 1 % (w/v) deoxycholate, 0.1% SDS, 150 mM NaCl, 50 mM NaF, 10 mM sodium orthovanadate, 1 mM EDTA, 1 mM DTT, and protease inhibitors (Complete™). Cells were subjected to three freeze-thaw cycles to ensure complete disruption. The cell debris was then removed by centrifugation
30 (13,000 g for 20 min at 4°C) and the supernatant incubated with 12 μ g of M2 anti-FLAG antibody (Sigma) overnight with mixing at 4°C. The immunocomplexes were then

collected with Protein A-Sepharose and washed four times with lysis buffer B. The immunoprecipitates were then subjected to SDS-PAGE on 12 % acrylamide gels, the gels dried and the phosphorylated hSK quantitated by Phosphorimager (Molecular Dynamics).

5 *In vitro phosphorylation of hSK1*

Purified recombinant hSK1 (1.5 µg) was incubated with recombinant ERK2 (0.1 µg; Upstate Biotechnology), 125 µM ATP and 5 µCi [γ^{32} P]ATP in 20 mM MOPS (pH 7.2) containing 20 mM MgCl₂, 25 mM β -glycerophosphate, 5 mM EGTA, 1 mM sodium
10 orthovanadate, and 1 mM DTT. The reaction was stopped after 20 min at 37°C by the addition of SDS-PAGE sample buffer, the mixture boiled for 5 min, subjected to SDS-PAGE on 12 % acrylamide gels, the gels dried, and the phosphorylated hSK1 quantitated by Phosphorimager.

15 *Sphingosine kinase assays*

SK activity was routinely determined using D-erythro-sphingosine (Biomol, Plymouth Meeting, PA) and [γ^{32} P]ATP (Geneworks, Adelaide, South Australia) as substrates, as described previously (Pitson *et al.* 2000). A unit (U) of SK activity is defined as the
20 amount of enzyme required to produce 1 pmol S1P/min.

EXAMPLE 2

Results

25 *hSK1 is phosphorylated in HEK293 cells in response to cell agonists*

The *in vivo* phosphorylation of hSK1 in HEK293 cells in response to TNF α and phorbol esters (PMA) was examined since phosphorylation is a common mechanism of regulating the catalytic activity of many eukaryotic. Metabolic labelling, with [32 P]orthophosphate, of
30 HEK293 cells stably over-expressing hSK1 1000-fold revealed that hSK1 is, indeed, phosphorylated, and that this phosphorylation increases rapidly in response to cell

exposure to TNF α (Fig 1) and PMA. Furthermore, this increase in hSK1 phosphorylation is highly correlated with the observed increase in sphingosine kinase activity in these cells in response to TNF α (Fig 1) and PMA. These observations suggested that phosphorylation plays a role in hSK1 activation. Further evidence for this was also obtained using a hSK1 mutant defective in its ability to bind TRAF2. This hSK1 mutant was is neither activated (Xia *et al.* 2002) or phosphorylated in response to TNF α (Fig 2). PMA, however, induces both activation (Xia *et al.* 2002) and phosphorylation of this mutant (Fig 2).

hSK1 is phosphorylated at Ser²²⁵

10

Analysis of the hSK1 sequence with the NetPhos phosphorylation site prediction program (Blom *et al.* 1999) allowed the identification of several possible phosphorylation sites in hSK1 that were also conserved in the mouse and monkey SK1 isoforms (Fig 3). Alanine mutagenesis of these possible sites revealed that the hSK1 mutant containing the Ser²²⁵ → Ala mutation was not phosphorylated (Fig 4). This suggested Ser²²⁵ as the sole physiologic phosphorylation site in hSK1. Further mutagenesis to alanine of Ser²²⁰ and Thr²²², two potentially phosphorylatable amino acids in the region around Ser²²⁵ had no effect on hSK1 phosphorylation (Fig 5). This indicated that the Ser²²⁵ → Ala mutation directly blocks phosphorylation at Ser²²⁵ rather than having a secondary effect on phosphorylation at an adjacent site (i.e. Ser²²⁰ or Thr²²²).

20

Activation of hSK1 requires phosphorylation at Ser²²⁵

The hSK^{S225A} mutant, when expressed in HEK293 cells, cannot be activated by cell treatment with TNF α or PMA (Fig 6). This provides strong evidence that phosphorylation Ser²²⁵ is essential for hSK1 activation.

25

ERK1, ERK2, and CDK2 specifically phosphorylate hSK1 at Ser²²⁵ in vitro

The amino acid sequence around Ser²²⁵ is SKTPAS²²⁵PVVVQ <400>3. The presence of a proline immediately C-terminal to Ser²²⁵ suggests a member of the proline-directed protein

30

kinase family is responsible for hSK1 phosphorylation. In particular, the PASP sequence of this region is reminiscent of the ERK1/2 substrate recognition motif, PxS/TP, where x represents a small, neutral amino acid (Chen *et al.* 2001). Experiments examining *in vitro* phosphorylation of purified recombinant hSK1 with ERK1, ERK2 and cyclin-dependent kinase 2 (CDK2) showed that all three kinases can phosphorylate hSK1 (Fig 7). Of these kinases, however, ERK2 showed the greatest efficiency of hSK1 phosphorylation. Further *in vitro* phosphorylation analysis with purified recombinant hSK^{S225A} mutant showed that ERK2 could not phosphorylate this protein (Fig 8). ERK2 was, however, still able to phosphorylate recombinant hSK1 mutants containing alanine mutations the only other potential proline-dependent kinase phosphorylation sites in hSK1 (Ser¹⁴⁸ and Thr²²²). This indicated that ERK2 specifically phosphorylates hSK1 at Ser²²⁵.

Increases in hSK1 phosphorylation in cells are blocked by a ERK1/2 pathway inhibitor, but not a CDK inhibitor

The *in vivo* phosphorylation of hSK1 in HEK293 cells in response to TNF α and PMA in the presence of PD98059 (a chemical inhibitor of the ERK1/2 pathway) and Olomoucine (a chemical inhibitor of CDKs) was examined. The results (Fig 9) show that PD98059 prevents TNF α - and PMA-induced increases in hSK1 phosphorylation. In contrast, the presence of Olomoucine does not block PMA-induced hSK1 phosphorylation, and actually appears to increase basal hSK1 phosphorylation in the absence of agonist stimulation. This data suggests the ERK1/2 pathway is necessary for hSK1 phosphorylation.

In vitro activation of hSK1 by ERK2

In vitro phosphorylation of purified recombinant hSK1 at Ser²²⁵ by ERK2 results in an approximate 4-fold increase in catalytic activity of hSK1 (Fig 10). This suggests that phosphorylation induces a conformational change in the protein structure of hSK1 that directly increases its catalytic efficiency.

Mutagenesis of Ser²²⁵ → Glu does not create a constitutively activated hSK1

By virtue of their negative charge the acidic amino acids glutamate and aspartate can sometimes mimic phosphorylated amino acids and create a protein conformation
 5 resembling the activated state. However, a hSK1 mutant containing the Ser²²⁵ → Glu mutation had similar activity to the wild-type hSK1 (Fig 11). Therefore, unlike the case in some other enzymes this mutation not create a constitutively activated hSK1.

Phospho-hSK1-specific polyclonal antibodies

10

The crude polyclonal antisera raised in rabbits against the phosphopeptide centred around Ser²²⁵ of hSK1 was highly reactive in ELISA towards the phosphopeptide, but also showed similar reactivity to the non-phosphorylated peptide (Fig 14). However, following affinity chromatography using the non-phosphorylated peptide to remove reactivity to this peptide,
 15 the antiserum showed high specificity in ELISA to the phosphopeptide (Fig 14). Furthermore, high specificity was also shown for the phosphorylated hSK1 protein in Western blots (Fig 15). In these Western blots, the antiserum showed reactivity towards ERK2 phosphorylated hSK1, but showed little reactivity towards the non-phosphorylated wildtype hSK1 (Fig 15) or the hSK^{S225A} mutant (Fig 16). Thus, we have used this purified
 20 antiserum to show TNFα and PMA-dependent increases in hSK1 phosphorylation (Fig 16).

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood
 25 that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

Table 3 Mutagenic oligonucleotides used for site-directed mutagenesis of hSK1

Mutation	Sequence
S148A	CGGCTGCTGGCGCCCATGAAC <400>4
S181A	TGTGGACCTCGAGGCTGAGAAGTA <400>5
Y184A	AGTGAGAAGGCTCGGCGCCTGGGGGAG <400>6
S220A	AAGAGTGGGCGCCAAGACAC <400>7
T222A	AAGAGTGGGATCCAAGGCGCCTGCCTCC <400>8
S225A	AAGACACCTGCGGCGCCCGTTGTG <400>9
S225E	ACACCTGCCGAACCGGTTGTGGTC <400>10
T250A	TCTCACTGGGCAGTGGTGC <400>11

BIBLIOGRAPHY

- Blom, N *et al.* (1999) *J Mol Biol* **294**, 1351-1362
Chen, Z *et al.* (2001) *Chem Rev* **101**, 2449-2476
Choi, OH *et al.* (1996) *Nature* **380**, 634-639
Cuvilliver, O *et al.* (1998) *J Biol Chem* **273**, 2910-2916
Desai, NN *et al.* (1992) *J Biol Chem* **267**, 23122-23128
Kleuser, B *et al.* (1998) *Cancer Res* **58**, 1817-1823
Mandala, SM *et al.* (2000) *Proc Natl Acad Sci USA* **97**, 7859-7964
Mattie, M *et al.* (1994) *J Biol Chem* **269**, 3181-3188
Melendez, A *et al.* (1998) *J Biol Chem* **273**, 9393-9402
Meyer zu Heringdorf, D *et al.* (1998) *EMBO J* **17**, 2830-2838
Meyer zu Heringdorf, D *et al.* (1999) *FEBS Lett* **461**, 217-221
Olivera, A & Spiegel, S. (1993) *Nature* **365**, 557-560
Pitson, SM *et al.* (2000) *Biochem J* **350**, 429-441
Pitson, SM *et al.* (2000) *J Biol Chem* **275**, 33945-33950
Rius, RA *et al.* (1997) *FEBS Lett* **417**, 173-176
Spiegel, S *et al.* (1998) *Ann N Y Acad Sci* **845**, 11-18
Su, Y *et al.* (1994) *J Biol Chem* **269**, 16512-16517
Van Veldhoven, PP *et al.* (2000) *Biochim Biophys Acta* **1487**, 128-134
Xia, P *et al.* (1998) *Proc Natl Acad Sci USA* **95**, 14196-14201
Xia, P *et al.* (2002) *J Biol Chem*, in press

DATED this 8th day of April, 2002

Medvet Science Pty Ltd
by its Patent Attorneys
DAVIES COLLISON CAVE

- 1 -

SEQUENCE LISTING

<110> Medvet Science Pty Ltd

<120> A Method of Modulating Cellular Activity - II

<130> 2519943/TDO

<160> 12

<170> PatentIn version 3.1

<210> 1

<211> 10

<212> PRT

<213> mammalian

<400> 1

Lys Thr Pro Ala Ser Pro Val Val Val Gln
1 5 10

<210> 2

<211> 14

<212> PRT

<213> mammalian

<400> 2

Cys Gly Ser Lys Thr Pro Ala Ser Pro Val Val Val Gln Gln
1 5 10

- 2 -

<210> 3
<211> 11
<212> PRT
<213> mammalian

<400> 3

Ser Lys Thr Pro Ala Ser Pro Val Val Val Gln
1 5 10

<210> 4
<211> 21
<212> DNA
<213> mammalian

<400> 4
cggctgctgg cgcccatgaa c

21

<210> 5
<211> 24
<212> DNA
<213> mammalian

<400> 5
tgtggacctc gaggctgaga agta

24

<210> 6
<211> 27
<212> DNA
<213> mammalian

<400> 6
agtgagaagg ctcggcgcct gggggag

27

- 3 -

<210> 7
<211> 20
<212> DNA
<213> mammalian

<400> 7
aagagtgggc gccaaagacac

20

<210> 8
<211> 28
<212> DNA
<213> mammalian

<400> 8
aagagtggga tccaaggcgc ctgcctcc

28

<210> 9
<211> 24
<212> DNA
<213> mammalian

<400> 9
aagacacctg cggcgcccgt tgtg

24

<210> 10
<211> 24
<212> DNA
<213> mammalian

<400> 10
acacctgccg aaccggttgt ggtc

24

- 4 -

<210> 11
 <211> 19
 <212> DNA
 <213> mammalian

<400> 11
 tctcactggg cagtgggtgc

19

<210> 12
 <211> 384
 <212> PRT
 <213> mammalian

<400> 12

Met Asp Pro Ala Gly Gly Pro Arg Gly Val Leu Pro Arg Pro Cys Arg
 1 5 10 15

Val Leu Val Leu Leu Asn Pro Arg Gly Gly Lys Gly Lys Ala Leu Gln
 20 25 30

Leu Phe Arg Ser His Val Gln Pro Leu Leu Ala Glu Ala Glu Ile Ser
 35 40 45

Phe Thr Leu Met Leu Thr Glu Arg Arg Asn His Ala Arg Glu Leu Val
 50 55 60

Arg Ser Glu Glu Leu Gly Arg Trp Asp Ala Leu Val Val Met Ser Gly
 65 70 75 80

Asp Gly Leu Met His Glu Val Val Asn Gly Leu Met Glu Arg Pro Asp
 85 90 95

Trp Glu Thr Ala Ile Gln Lys Pro Leu Cys Ser Leu Pro Ala Gly Ser
 100 105 110

- 5 -

Gly Asn Ala Leu Ala Ala Ser Leu Asn His Tyr Ala Gly Tyr Glu Gln
115 120 125

Val Thr Asn Glu Asp Leu Leu Thr Asn Cys Thr Leu Leu Cys Arg
130 135 140

Arg Leu Leu Ser Pro Met Asn Leu Leu Ser Leu His Thr Ala Ser Gly
145 150 155 160

Leu Arg Leu Phe Ser Val Leu Ser Leu Ala Trp Gly Phe Ile Ala Asp
165 170 175

Val Asp Leu Glu Ser Glu Lys Tyr Arg Arg Leu Gly Glu Met Arg Phe
180 185 190

Thr Leu Gly Thr Phe Leu Arg Leu Ala Ala Leu Arg Thr Tyr Arg Gly
195 200 205

Arg Leu Ala Tyr Leu Pro Val Gly Arg Val Gly Ser Lys Thr Pro Ala
210 215 220

Ser Pro Val Val Val Gln Gln Gly Pro Val Asp Ala His Leu Val Pro
225 230 235 240

Leu Glu Glu Pro Val Pro Ser His Trp Thr Val Val Pro Asp Glu Asp
245 250 255

Phe Val Leu Val Leu Ala Leu Leu His Ser His Leu Gly Ser Glu Met
260 265 270

Phe Ala Ala Pro Met Gly Arg Cys Ala Ala Gly Val Met His Leu Phe
275 280 285

Tyr Val Arg Ala Gly Val Ser Arg Ala Met Leu Leu Arg Leu Phe Leu
290 295 300

Ala Met Glu Lys Gly Arg His Met Glu Tyr Glu Cys Pro Tyr Leu Val
305 310 315 320

- 6 -

Tyr Val Pro Val Val Ala Phe Arg Leu Glu Pro Lys Asp Gly Lys Gly
325 330 335

Met Phe Ala Val Asp Gly Glu Leu Met Val Ser Glu Ala Val Gln Gly
340 345 350

Gln Val His Pro Asn Tyr Phe Trp Met Val Ser Gly Cys Val Glu Pro
355 360 365

Pro Pro Ser Trp Lys Pro Gln Gln Met Pro Pro Pro Glu Glu Pro Leu
370 375 380

Figure 1




Figure 2




Figure 3

1 MDPAGGPRGVLP RPCRVLVLLNPRGGKGKALQLFRSHVQPLLAAEAEISFTLMLTERRNHARELVRSEELG 70

71 RWDALVVMMSGDGLMHEVVNGLMERPDWETAIQKPLCSLPAGSGNALAASLNHYAGYEQVTNEDLLTNCTL 140

141 LLCRRLLSPMNLISLHTASGLRLLFSVLSLAWGFIADVDDLSPKYYRRLGEMRFTLTGTFRLAALRTYRGRL 210

211 AYLPGVRVGSKTPASPVVVQQGPVDAHLVPLEEPVPSHNTTVDPDEDFVLVALLHSHLGSEMFAAPMGRC 280

281 AAGVMHLFYVRAGVSRAMLLRLFLAMEKGRHMEYECPYLVVVPVAFRLEPKDGKGMFAVDGELMVSEAV 350

351 QGQVHPNIFYFMVSGCVEPPPSWKPPQMPPEEPL 384

Figure 4

4/15

pcDNA3	WT	S148A	S181A	Y184A	S225A	T250A
-	-	+	+	-	+	+
+	-	+	+	-	+	+

PMA

³²P incorporation

Protein




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15

